

REPRESENTATIVE PUBLICATIONS BY MMI FACULTY DIRECTORS AND CONSULTANTS

MMI's radiologists and imaging scientists have published extensively on techniques for knee imaging and cartilage evaluation. Sample publications are featured below for reference. Similar bibliographies exist for joint recon, trauma, fracture repair, etc.

- Mosher TJ. Degenerative Disease: Cartilage Anatomy, Physiology and Advanced Imaging. *Imaging of the Musculoskeletal System*. Thomas Pope ed. Philadelphia, Saunders Elsevier; 2008
- Gold G, Chen CA, Koo S, Hargreaves BA, Bangerter NK. Recent advances in MRI of articular cartilage. *Am J Roentgenol*. 2009;193(3):628-38.
- Crema MD, Roemer FW, Marra MD, Burstein D, Gold GE, Eckstein F, Baum T, Mosher TJ, Carrino JA, Guermazi A. Articular cartilage in the knee: current MR imaging techniques and applications in clinical practice and research. *Radiographics*; 2011 31:37-62.
- Mosher TJ, Zhang Z, Reddy R, Boudhar S, Milestone B, Morrison W, Kwok K, Eckstein F, Witschey W, Borthakur A. Knee articular cartilage damage in osteoarthritis: analysis of MR image biomarker reproducibility in ACRIN-PA 4001 multicenter trial. *Radiology*; 2011 258:832-842.
- Trattnig S, Domayer S, Welsch GW, Mosher TJ, Eckstein F. MR imaging of cartilage and its repair in the knee--a review. *Eur Radiol* 2009; 19:1582-1594.
- Ly JQ, Beall DP. MR Imaging of the Painful Knee: Correlation with Conventional Radiography *Appl Radiol* 2003;32:39-45
- Beall DP, Scott WW, Eng J, Johnson CA. Improved Demonstration of Cartilage Narrowing in the Knee Joint Using Standing PA Flexed Radiographs [Abstract]. *Radiology* 1999;213(P)(Suppl):203.
- Gold GE, Burstein D, Dardzinski B, Lang P, Boada F, Mosher TJ. MRI of articular cartilage in OA: novel pulse sequences and compositional/functional markers. *Osteoarthritis Cartilage* 2006; 14 Suppl A:A76-86.
- Eckstein F, Ateshian G, Burgkart R, Burstein D, Cicuttini F, Dardzinski B, Gray M, Link TM, Majumdar S, Mosher TJ, Peterfy C, Totterman S, Waterton J, Winalski CS, Felson D. Proposal for a nomenclature for Magnetic Resonance Imaging based measures of articular cartilage in osteoarthritis. *Osteoarthritis Cartilage* 2006; 14:974-983.
- Choi JA, Gold G. MR imaging of articular cartilage physiology. *Magn Reson Imaging Clin N Am*. 2011;19(2):249-82.
- Chen CA, Kijowski R, Shapiro LM, Tuite MJ, Davis KW, Klaers JL, Block WF, Reeder SB, Gold G. Cartilage morphology at 3.0T: assessment of three-dimensional magnetic resonance imaging techniques. *J MRI*. 2010;32:173-83.
- Friedrich KM, Reiter G, Kaiser B, Mayerhöfer M, Deimling M, Jellus V, Horger W, Trattnig S, Schweitzer ME, Salomonowitz E. High-resolution cartilage imaging of the knee at 3T: basic evaluation of modern isotropic 3D MR-sequences. *Eur J Radiol*. 2011 Jun;78(3):398-405.
- Gold G, McCauley TR, Gray ML, Disler DG. What's new in cartilage? *Radiographics*. 2003 Sep-Oct;23(5):1227-42.
- Carrino JA, Blum M, Parellada JA, Schweitzer ME, Morrison WB. MRI of bone marrow edema-like signal in the pathogenesis of subchondral cysts. *Osteoarthritis Cartilage*. Oct; 14(10):1081-5.
- Ling W, Regatte RR, Schweitzer ME, Jerschow A. Behavior of ordered sodium in enzymatically depleted cartilage tissue. *Magn Reson Med*. 2006 Nov; 56(5):1151-5.
- Pakin SK, Schweitzer ME, Regatte RR. 3D-T1rho quantitation of patellar cartilage at 3.0T. *J Magn Reson Imaging*. 2006 Dec;24(6): 1357-63.
- Ling W, Regatte RR, Schweitzer ME, Jerschow A. Characterization of bovine patellar cartilage. *NMR Biomedicine* 2008;21:289-295.
- Wang L, Schweitzer ME, Pauda A Regatte RR. Rapid 3D-T1 Mapping of Cartilage with Variable-Flip-Angle (VFA) and Parallel Imaging at 3.0T. *JMRI* 2007; 27:1171-5
- Welsch GH, Mamisch TC, Hughes T, Zilkens C, Quirbach S, Scheffler K, Draff O, Schweitzer ME, Szomolanyi P, Trattnig S. In vivo biochemical 7.0 Tesla magnetic resonance: preliminary results of dGEMRIC, zonal T2, and T2* mapping of articular cartilage. *Invest Radiol*. 2008 Sept;43(9):619-26.
- Friedrich KM, Shepard T, de Oliveira VS, Wang L, Babb JS, Schweitzer ME, Regatte R. T2 measurements of cartilage in osteoarthritis patients with meniscal tears. *AJR* 2009 Nov; 193:W411-5.
- Vives MJ, Homesley D, Cicotti MG, Schweitzer ME. Evaluation of recurring meniscal tears with gadolinium-enhanced magnetic resonance imaging: a randomized, prospective study. *Am J Sports Med*. 203(6):868-73.
- White LW, Schweitzer ME, Johnson WJ, Amster BJ, Oliveri MP, Russell K. The role of T2-weighted fast-spin-echo imaging in the diagnosis of meniscal tears. *JMRI* 1996; 6:874-877.
- White LW, Schweitzer ME, Morrison WB, Deely DM. The effect of training and experience on the magnetic resonance imaging interpretation of meniscal tears. *Arthroscopy* 1997; 13:224-228.

- Blair TR, Schweitzer ME, Resnick D. Meniscal cysts causing bone erosion: retrospective analysis of seven cases. *Clin Imaging* 1999; 23:134-138.
- Hodler J, Berthiaume M, Schweitzer ME, et al. Knee joint hyaline cartilage defects: A comparative study of MR and anatomic sections. *J Comput Assist Tomogr* 1992; 16:597-680.
- Lektrakul N, Skaf A, Yeh L, Roger B, Schweitzer ME, Blasbalg R, Resnick D. Pericruciate meniscal cysts arising from tears of the posterior horn of the medial meniscus: MR imaging features that simulate posterior cruciate ganglion cysts. *AJR* 1999; 172:1575-1579.
- White LM, Schweitzer ME, Weishaupt D, Kramer J, Davis A, Marks PH. Diagnosis of recurrent meniscal tears: Prospective evaluation of conventional MR imaging, indirect MR arthrography, and direct MR arthrography. *Radiology* 2002
- Lim P, Schweitzer ME, Bhatia M, Giuliano V, Kaneriya PP, Senyk RM, Oliveri M, Johnson W, Amster B, Parker L. Repeat tear of postoperative meniscus: potential MR imaging signs. *Radiology* 1999; 210:183-188.
- Vance K, Meredick R, Schweitzer ME, Lubowitz JH. Magnetic resonance imaging of the postoperative meniscus. *Arthroscopy*. 2009 May;25(5):522-30.
- Rosenthal P, Abellana V, Greenberg J, Schweitzer ME, Abramson SB. Synovial but not cartilage volumes on MRI predict radiographic severity of knee OA. *Osteoarthritis and Cartilage*, 2007.15,Suppl.3:(C29).
- Krasnokutsky S, Samuels J, Attur M, Regatte R, Babb J, Rosenthal P, Abellana V, Greenberg J, Schweitzer ME, Abramson SB. Synovial but not cartilage volumes on MRI predict radiographic severity of knee OA. *Arthritis Rheum* 2008
- Friedrich KM, Shepard T, Chang G, Wang L, Babb JS, Schweitzer ME, Regatte R. Does joint alignment affect the T2 values of cartilage in patients with knee osteoarthritis? *Eur Radiol*. 2010;20(6):1532-8.
- van Leersum M, Schweitzer ME, Gannon F, Vinitski S, Finkel G, Mitchell DG. Thickness of patellofemoral articular cartilage as measured on MR imaging: Sequence comparison of accuracy, reproducibility, and interobserver variation. *Skeletal Radiol* 1995; 24:431-435.
- Pakin SK, Schweitzer ME, Regatte RR. 3D-T1rho quantitation of patellar cartilage at 3.0T. *J Magn Reson Imaging*. 2006 Dec;24(6): 1357-63.
- Ling W, Regatte RR, Schweitzer ME, Jerschow A. Characterization of bovine patellar cartilage. *NMR Biomedicine* 2008;21:289-295.
- Gold G, Hargreaves BA, Vasanawala SS, Webb JD, Shimakawa AS, Brittain JH, Beaulieu CF. Articular cartilage of the knee: evaluation with fluctuating equilibrium MR imaging--initial experience in healthy volunteers. *Radiology*. 2006;238(2):712-8.
- Gold G, Reeder SB, Yu H, Kornaat P, Shimakawa AS, Johnson JW, Pelc NJ, Beaulieu CF, Brittain JH. Articular cartilage of the knee: rapid three-dimensional MR imaging at 3.0 T with IDEAL balanced steady-state free precession--initial experience. *Radiology*. 2006;240(2):546-51.
- Miller KL, Hargreaves BA, Gold G, Pauly JM. Steady-state diffusion-weighted imaging of in vivo knee cartilage. *Magn Reson Med*. 2004;51(2):394-8.
- Vasnavala SS, Pauly JM, Nishimura DG, Gold G. MR imaging of knee cartilage with FEMR. *Skeletal Radiol*. 2002;31(10):574-80.
- Gold G, Thedens DR, Pauly JM, Fechner KP, Bergman G, Beaulieu CF, Macovski A. MR imaging of articular cartilage of the knee: New methods using ultrashort TEs. *AJR Am J Roentgenol*. 1998;170(5):1223-6.
- Gold G, Bergman AG, Pauly JM, Lang P, Butts RK, Beaulieu CF, Hargreaves B, Frank L, Boutin RD, Macovski A, Resnick D. Magnetic resonance imaging of knee cartilage repair. *Top Magn Reson Imaging*. 1998;9(6):377-92.
- Gold GE, Reeder SB, Beaulieu CF. Advanced MR imaging of the shoulder: dedicated cartilage techniques. *Magn Reson Imaging Clin N Am*. 2004;12(1):143-59, vii.
- Staroswiecki E, Granlund KL, Alley MT, Gold G, Hargreaves BA. Simultaneous estimation of T(2) and apparent diffusion coefficient in human articular cartilage in vivo with a modified three-dimensional double echo steady state (DESS) sequence at 3 T. *Magn Reson Med*. 2012;67(4):1086-96.
- Stikov N, Keenan KE, Pauly JM, Smith RL, Dougherty RF, Gold G. Cross-relaxation imaging of human articular cartilage. *Magn Reson Med*. 2011;66(3):725-34.
- Rakow-Penner R, Gold G, Daniel B, Stevens K, Rosenberg J, Mazin S, Pauly J, Glover GH. Reduction of truncation artifacts in rapid 3D articular cartilage imaging. *J Magn Reson Imaging*. 2008;27(4):860-5.
- Gold G, Hargreaves BA, Stevens KJ, Beaulieu CF. Advanced magnetic resonance imaging of articular cartilage. *Orthop Clin North Am*. 2006;37(3):331-47, vi.

- Gold G, Burstein D, Dardzinski B, Lang P, Boada F, Mosher T. MRI of articular cartilage in OA: novel pulse sequences and compositional/functional markers. *Osteoarthritis Cartilage*. 2006;14 Suppl A:A76-86.
- Gold G, Hargreaves BA, Reeder SB, Vasanawala SS, Beaulieu CF. Controversies in protocol selection in the imaging of articular cartilage. *Semin Musculoskelet Radiol*. 2005;9(2):161-72.
- Gold G, Fuller SE, Hargreaves BA, Stevens KJ, Beaulieu CF. Driven equilibrium magnetic resonance imaging of articular cartilage: initial clinical experience. *J Magn Reson Imaging*. 2005;21(4):476-81.
- Kornaat PR, Reeder SB, Koo S, Brittain JH, Yu H, Andriacchi TP, Gold G. MR imaging of articular cartilage at 1.5T and 3.0T: comparison of SPGR and SSFP sequences. *Osteoarthritis Cartilage*. 2005;13(4):338-44.
- Reeder SB, Pelc NJ, Alley MT, Gold G. Rapid MR imaging of articular cartilage with steady-state free precession and multipoint fat-water separation. *AJR Am J Roentgenol*. 2003;180(2):357-62.
- Gold G, Beaulieu CF. Future of MR imaging of articular cartilage. *Semin Musculoskelet Radiol*. 2001;5(4):313-27.
- Hargreaves BA, Gold G, Lang PK, Conolly SM, Pauly JM, Bergman G, Vandevenne J, Nishimura DG. MR imaging of articular cartilage using driven equilibrium. *Magn Reson Med*. 1999;42(4):695-703.
- Keenan KE, Besier TF, Pauly JM, Han E, Rosenberg J, Smith RL, Delp SL, Beaupre GS, Gold G. Prediction of glycosaminoglycan content in human cartilage by age, T1# and T2 MRI. *Osteoarthritis Cartilage*. 2011;19(2):171-9.
- McWalter EJ, Gold G. UTE T2# mapping detects sub-clinical meniscus degeneration. *Osteoarthritis Cartilage*. 2012;20(6):471-2.
- Chen CA, Lu W, John CT, Hargreaves BA, Reeder SB, Delp SL, Siston RA, Gold G. Multiecho IDEAL gradient-echo water-fat separation for rapid assessment of cartilage volume at 1.5 T: initial experience. *Radiology*. 2009;252(2):561-7.
- Keenan KE, Kourtis LC, Besier TF, Lindsey DP, Gold G, Delp SL, Beaupre GS. New resource for the computation of cartilage biphasic material properties with the interpolant response surface method. *Comput Methods Biomech Biomed Engin*. 2009;12(4):415-22.
- Kornaat PR, Koo S, Andriacchi TP, Bloem JL, Gold G. Comparison of quantitative cartilage measurements acquired on two 3.0T MRI systems from different manufacturers. *J Magn Reson Imaging*. 2006;23(5):770-3.
- Koo S, Gold G, Andriacchi TP. Considerations in measuring cartilage thickness using MRI: factors influencing reproducibility and accuracy. *Osteoarthritis Cartilage*. 2005;13(9):782-9.
- Hargreaves BA, Gold G, Beaulieu CF, Vasanawala SS, Nishimura DG, Pauly JM. Comparison of new sequences for high-resolution cartilage imaging. *Magn Reson Med*. 2003;49(4):700-9.
- Lazovic-Stojkovic J, Mosher TJ, Smith HE, Yang QX, Dardzinski BJ, Smith MB. Interphalangeal Joint Cartilage: High-Spatial-Resolution in Vivo MR T2 Mapping--A Feasibility Study. *Radiology* 2004; 233:292-296.
- Smith HE, Mosher TJ, Dardzinski BJ, Collins BG, Collins CM, Yang QX, Schmithorst VJ, Smith MB: Spatial Variation in Cartilage T2 of the Knee. *J Magn Reson Imag*. 2001;14:50-55
- Mosher TJ, Smith HE, Collins C, Liu Y, Hancy J, Dardzinski BJ, Smith MB. Change in knee cartilage T2 at MR imaging after running: a feasibility study. *Radiology* 2005;234(1):245-9.
- Mosher TJ, Liu Y, Torok CM. Functional cartilage MRI T2 mapping: evaluating the effect of age and training on knee cartilage response to running. *Osteoarthritis Cartilage*; 2010 18:358-364.
- Mosher TJ, Dardzinski BJ. Cartilage MRI T2 relaxation time mapping: overview and applications. *Semin Musculoskelet Radiol* 2004;8(4):355-68.
- Dardzinski BJ, Mosher TJ, Li S, Van Slyke MA, Smith MB: Spatial Variation of T2 in Human Articular Cartilage. *Radiology*, 205(2):546-550, 1997.
- Mosher TJ, Pruett SW: Magnetic Resonance Imaging of Superficial Cartilage Lesions: Role of Contrast in Lesion Detection. *J Magn. Reson. Imag*. 10:178-182, 1999.
- Mosher TJ, Dardzinski BJ, Smith MB: Human Articular Cartilage: Influence of Aging and Early Symptomatic Degeneration on the Spatial Variation of T2 – Preliminary Findings at 3T. *Radiology*, 214(1):259-266, 2000.
- Mosher TJ, Smith H, Dardzinski BJ, Schmithorst VJ, Smith MB: MR Imaging and T2 Mapping of Femoral Cartilage: In Vivo Determination of the Magic Angle Effect. *AJR* 177:665-669, 2001.
- Mosher TJ, Chen Q, Smith MB: 1H Magnetic Resonance Spectroscopy of Nanomelic Chicken Cartilage: Effect of Aggrecan Depletion on Cartilage T2. *Osteoarthritis and Cartilage* 11(10):709-715, 2003.
- Mosher TJ, Collins CM, Smith HE, Moser LE, Sivarajah RT, Dardzinski BJ, Smith MB: Effect of gender on in vivo cartilage magnetic resonance imaging T2 mapping. *J Magn Reson Imaging* 2004; 19:323-328
- Mosher, TJ, Liu, Y, Yang, QX, Yao J, Smith R, Dardzinski, BJ, Smith MB: Age dependency of cartilage magnetic resonance imaging T2 relaxation times in asymptomatic women. *Arthritis Rheum* 2004; 50:2820.